



# CHAPTER 1

*Sequences And Series*

---

# CONTENTS

<b>1</b>	<b>Arithmetic sequences</b>	<b>1</b>
1.1	Arithmetic sequences . . . . .	1
<b>2</b>	<b>Geometric sequences</b>	<b>10</b>
<b>3</b>	<b>Series</b>	<b>15</b>
3.1	Series . . . . .	15
3.2	Sigma notation . . . . .	16
<b>4</b>	<b>Finite arithmetic series</b>	<b>20</b>
4.1	Finite arithmetic series . . . . .	20
4.2	General formula for a finite arithmetic series . . . . .	21
<b>5</b>	<b>Finite geometric series</b>	<b>27</b>
5.1	Finite geometric series . . . . .	27
5.2	General formula for a finite geometric series . . . . .	28
<b>6</b>	<b>Infinite series</b>	<b>32</b>
6.1	Infinite series . . . . .	32
6.2	Infinite geometric series . . . . .	35
<b>7</b>	<b>Summary</b>	<b>38</b>
<b>8</b>	<b>Exercises</b>	<b>40</b>
8.1	Exercise 1 . . . . .	40
8.2	Exercise 2 . . . . .	42
8.3	Exercise 3 . . . . .	43
8.4	Exercise 4 . . . . .	45
8.5	Exercise 5 . . . . .	46
8.6	Exercise 6 . . . . .	48
8.7	Exercise 7 . . . . .	49
<b>9</b>	<b>Answers for Exercises</b>	<b>50</b>
9.1	Exercise 1 . . . . .	50
9.2	Exercise 2 . . . . .	53
9.3	Exercise 3 . . . . .	53
9.4	Exercise 4 . . . . .	54
9.5	Exercise 5 . . . . .	55

---

9.6	Exercise 6	56
9.7	Exercise 7	57

August 26, 2021

---

# 1 ARITHMETIC SEQUENCES

In earlier grades we learnt about number patterns, which included linear sequences with a common difference and quadratic sequences with a common second difference. We also looked at completing a sequence and how to determine the general term of a sequence.

In this chapter we also look at geometric sequences, which have a constant ratio between consecutive terms. We will learn about arithmetic and geometric series, which are the summing of the terms in sequences.

## 1.1 Arithmetic sequences

An arithmetic sequence is a sequence where consecutive terms are calculated by adding a constant value (positive or negative) to the previous term. We call this constant value the common difference ( $d$ ).

For example,

$$3; 0; -3; -6; -9; \dots$$

This is an arithmetic sequence because we add  $-3$  to each term to get the next term:

First term	$T_1$		3
Second term	$T_2$	$3 + (-3) =$	0
Third term	$T_3$	$0 + (-3) =$	-3
Fourth term	$T_4$	$-3 + (-3) =$	-6
Fifth term	$T_5$	$-6 + (-3) =$	-9
$\vdots$	$\vdots$	$\vdots$	$\vdots$

### The general term for an arithmetic sequence

For a general arithmetic sequence with first term  $a$  and a common difference  $d$ , we can generate the following terms:

$$T_1 = a$$

$$T_2 = T_1 + d = a + d$$

$$T_3 = T_2 + d = (a + d) + d = a + 2d$$

$$T_4 = T_3 + d = (a + 2d) + d = a + 3d$$

$$\vdots \quad \quad \quad \vdots \quad \quad \quad \vdots \quad \quad \quad \vdots$$

$$T_n = T_{n-1} + d = (a + (n-2)d) + d = a + (n-1)d$$

Therefore, the general formula for the  $n^{\text{th}}$  term of an arithmetic sequence is:

$$T_n = a + (n-1)d$$

### Definition: Arithmetic sequence

An arithmetic (or linear) sequence is an ordered set of numbers (called terms) in which each new term is calculated by adding a constant value to the previous term:

$$T_n = a + (n - 1)d$$

where

- $T_n$  is the  $n^{\text{th}}$  term;
- $n$  is the position of the term in the sequence;
- $a$  is the first term;
- $d$  is the common difference.

### Test for an arithmetic sequence

To test whether a sequence is an arithmetic sequence or not, check if the difference between any two consecutive terms is constant:

$$d = T_2 - T_1 = T_3 - T_2 = \dots = T_n - T_{n-1}$$

If this is not true, then the sequence is not an arithmetic sequence.

### WORKED EXAMPLE 1: ARITHMETIC SEQUENCE

#### QUESTION

Given the sequence  $-15; -11; -7; \dots 173$ .

1. Is this an arithmetic sequence?
2. Find the formula of the general term.
3. Determine the number of terms in the sequence.

#### SOLUTION

##### Step 1: Check if there is a common difference between successive terms

$$T_2 - T_1 = -11 - (-15) = 4$$

$$T_3 - T_2 = -7 - (-11) = 4$$

$\therefore$  This is an arithmetic sequence with  $d = 4$

##### Step 2: Determine the formula for the general term


Write down the formula and the known values:

$$T_n = a + (n - 1)d$$

### WORKED EXAMPLE 1: ARITHMETIC SEQUENCE (continued)

$$a = -15; \quad d = 4$$

$$\begin{aligned} T_n &= a + (n - 1)d \\ &= -15 + (n - 1)(4) \\ &= -15 + 4n - 4 \\ &= 4n - 19 \end{aligned}$$



### WORKED EXAMPLE 1: ARITHMETIC SEQUENCE (continued)

A graph was not required for this question but it has been included to show that the points of the arithmetic sequence lie in a straight line.

Note: The numbers of the sequence are natural numbers ( $n \in \{1; 2; 3; \dots\}$ ) and therefore we should not connect the plotted points. In the diagram above, a dotted line has been used to show that the graph of the sequence lies on a straight line.

#### Step 3: Determine the number of terms in the sequence

$$T_n = a + (n - 1)d$$

$$173 = 4n - 19$$

$$192 = 4n$$

$$\therefore n = \frac{192}{4}$$

$$= 48$$

$$\therefore T_{48} = 173$$

#### Step 4: Write the final answer

Therefore, there are 48 terms in the sequence.

### Arithmetic mean

The arithmetic mean between two numbers is the number half-way between the two numbers. In other words, it is the average of the two numbers. The arithmetic mean and the two terms form an arithmetic sequence.

For example, the arithmetic mean between 7 and 17 is calculated:


$$\begin{aligned}\text{Arithmetic mean} &= \frac{7 + 17}{2} \\ &= 12\end{aligned}$$

$\therefore 7; 12; 17$  is an arithmetic sequence

$$T_2 - T_1 = 12 - 7 = 5$$

$$T_3 - T_2 = 17 - 12 = 5$$

Plotting a graph of the terms of a sequence sometimes helps in determining the type of sequence involved. For an arithmetic sequence, plotting  $T_n$  vs.  $n$  results in the following graph:



- If the sequence is arithmetic, the plotted points will lie in a straight line.
- Arithmetic sequences are also called linear sequences, where the common difference ( $d$ ) is the gradient of the straight line.

$$T_n = a + (n - 1)d$$

can be written as  $T_n = d(n - 1) + a$

which is of the same form as  $y = mx + c$




**Definition: Quadratic sequence**

A quadratic sequence is a sequence of numbers in which the second difference between any two consecutive terms is constant.

The general formula for the  $n^{\text{th}}$  term of a quadratic sequence is:

$$T_n = an^2 + bn + c$$




It is important to note that the first differences of a quadratic sequence form an arithmetic sequence. This sequence has a common difference of  $2a$  between consecutive terms. In other words, a linear sequence results from taking the first differences of a quadratic sequence.

**WORKED EXAMPLE 2: QUADRATIC SEQUENCE**

**QUESTION**

Consider the pattern of white and blue blocks in the diagram below.

- Determine the sequence formed by the white blocks ( $w$ ).
- Find the sequence formed by the blue blocks ( $b$ ).



Pattern number (n)	1	2	3	4	5	6	n
No. of white blocks (w)							
Common difference (d)							

Pattern number (n)	1	2	3	4	5	6	$n$
No. of blue blocks (b)							
Common difference (d)							

### WORKED EXAMPLE 2: QUADRATIC SEQUENCE (continued)

#### SOLUTION

**Step 1: Use the diagram to complete the table for the white blocks**

Pattern number (n)	1	2	3	4	5	6	$n$
No. of white blocks (w)	4	8	12	16	20	24	$4n$
Common difference (d)		4	4	4	4	4	

We see that the next term in the sequence is obtained by adding 4 to the previous term, therefore the sequence is linear and the common difference ( $d$ ) is 4.

The general term is:

$$\begin{aligned}
 T_n &= a + (n - 1)d \\
 &= 4 + (n - 1)(4) \\
 &= 4 + 4n - 4 \\
 &= 4n
 \end{aligned}$$

**Step 2: Use the diagram to complete the table for the blue blocks**

Pattern number (n)	1	2	3	4	5	6
No. of blue blocks (b)	0	1	4	9	16	24
difference (d)		1	3	5	7	9

We notice that there is no common difference between successive terms. However, there is a pattern and on further investigation we see that this is in fact a quadratic sequence:


Pattern number (n)	1	2	3	4	5	6	$n$
No. of blue blocks (b)	0	1	4	9	16	25	$(n - 1)^2$
First difference	—	1	3	5	7	9	—
Second difference	—	—	2	2	2	2	—
Pattern	$(1 - 1)^2$	$(2 - 1)^2$	$(3 - 1)^2$	$(4 - 1)^2$	$(5 - 1)^2$	$(6 - 1)^2$	$(n - 1)^2$

$$T_n = (n - 1)^2$$

**Step 3: Draw a graph of  $T_n$  vs.  $n$  for each sequence**

White blocks:  $T_n = 4n$

Blue blocks:  $T_n = (n - 1)^2$   
 $= n^2 - 2n + 1$



Since the numbers of the sequences are natural numbers ( $n \in \{1; 2; 3; \dots\}$ ) we should not connect the plotted points. In the diagram above, a dotted line has been used to show that the graph of the sequence

---

formed by the white blocks ( $w$ ) is a straight line and the graph of the sequence formed by the blue blocks ( $b$ ) is a parabola.

## 2 GEOMETRIC SEQUENCES

### Definition: Geometric sequence

A geometric sequence is a sequence of numbers in which each new term (except for the first term) is calculated by multiplying the previous term by a constant value called the constant ratio ( $r$ ).


This means that the ratio between consecutive numbers in a geometric sequence is a constant (positive or negative). We will explain what we mean by ratio after looking at the following example.

### Example: A flu epidemic

#### Example: A flu epidemic

Influenza (commonly called “flu”) is caused by the influenza virus, which infects the respiratory tract (nose, throat, lungs). It can cause mild to severe illness that most of us get during winter time. The influenza virus is spread from person to person in respiratory droplets of coughs and sneezes. This is called “droplet spread”. This can happen when droplets from a cough or sneeze of an infected person are propelled through the air and deposited on the mouth or nose of people nearby. It is good practice to cover your mouth when you cough or sneeze so as not to infect others around you when you have the flu. Regular hand washing is an effective way to prevent the spread of infection and illness.

Assume that you have the flu virus, and you forgot to cover your mouth when two friends came to visit while you were sick in bed. They leave, and the next day they also have the flu. Let’s assume that each friend in turn spreads the virus to two of their friends by the same droplet spread the following day. Assuming this pattern continues and each sick person infects 2 other friends, we can represent these events in the following manner:



### Example: A flu epidemic (continued)

Each person infects two more people with the flu virus.

We can tabulate the events and formulate an equation for the general case:

Day (n)	No. of newly-infected people
1	$2 = 2$
2	$4 = 2 \times 2 = 2 \times 2^1$
3	$8 = 2 \times 4 = 2 \times 2 \times 2 = 2 \times 2^2$
4	$16 = 2 \times 8 = 2 \times 2 \times 2 \times 2 = 2 \times 2^3$
5	$32 = 2 \times 16 = 2 \times 2 \times 2 \times 2 \times 2 = 2 \times 2^4$
$\vdots$	$\vdots$
$n$	$2 \times 2 \times 2 \times 2 \times \dots \times 2 = 2 \times 2^{n-1}$

The above table represents the number of **newly-infected** people after  $n$  days since you first infected your 2 friends.

You sneeze and the virus is carried over to 2 people who start the chain. ( $a = 2$ ) The next day, each one then infects 2 of their friends. Now 4 people are newly-infected. Each of them infects 2 people the third day, and 8 new people are infected, and so on. These events can be written as a geometric sequence:

$$2; 4; 8; 16; 32; \dots$$

Note the constant ratio ( $r = 2$ ) between the events. Recall from the linear arithmetic sequence how the common difference between terms was established. In the geometric sequence we can determine the constant ratio ( $r$ ) from:

$$\frac{T_2}{T_1} = \frac{T_3}{T_2} = r$$

More generally,

$$\frac{T_n}{T_{n-1}} = r$$

### The general term for a geometric sequence

From the flu example above we know that  $T_1 = 2$  and  $r = 2$  and we have seen from the table that the  $n^{\text{th}}$  term is given by  $T_n = 2 \times 2^{n-1}$ .

The general geometric sequence can be expressed as:

$$\begin{aligned}T_1 &= a &&= ar^0 \\T_2 &= a \times r &&= ar^1 \\T_3 &= a \times r \times r &&= ar^2 \\T_4 &= a \times r \times r \times r &&= ar^3 \\T_n &= a \times [r \times r \dots (n-1) \text{ times}] &&= ar^{n-1}\end{aligned}$$

Therefore the general formula for a geometric sequence is:

$$T_n = ar^{n-1}$$

where

- $a$  is the first term in the sequence;
- $r$  is the constant ratio.

#### Test for a geometric sequence

To test whether a sequence is a geometric sequence or not, check if the ratio between any two consecutive terms is constant:

$$\frac{T_2}{T_1} = \frac{T_3}{T_2} = \frac{T_n}{T_{n-1}} = r$$

If this condition does not hold, then the sequence is not a geometric sequence.

#### WORKED EXAMPLE 3: FLU EPIDEMIC

##### QUESTION

We continue with the previous flu example, where  $T_n$  is the number of newly-infected people after  $n$  days:

$$T_n = 2 \times 2^{n-1}$$

1. Calculate how many newly-infected people there are on the tenth day.
2. On which day will 16 384 people be newly-infected?

##### SOLUTION

**Step 1: Write down the known values and the general formula**

$$a = 2$$

$$r = 2$$

$$T_n = 2 \times 2^{n-1}$$

### WORKED EXAMPLE 3: FLU EPIDEMIC (continued)

**Step 2: Use the general formula to calculate  $T_{10}$**

Substitute  $n = 10$  into the general formula:

$$\begin{aligned}T_n &= a \times r^{n-1} \\ \therefore T_{10} &= 2 \times 2^{10-1} \\ &= 2 \times 2^9 \\ &= 2 \times 512 \\ &= 1024\end{aligned}$$

On the tenth day, there are 1 024 newly-infected people.

**Step 3: Use the general formula to calculate  $n$**

We know that  $T_n = 16\,384$  and can use the general formula to calculate the corresponding value of  $n$ :

$$\begin{aligned}T_n &= ar^{n-1} \\ 16\,384 &= 2 \times 2^{n-1} \\ \frac{16\,384}{2} &= 2^{n-1} \\ 8\,192 &= 2^{n-1}\end{aligned}$$

We can write 8 192 as  $2^{13}$

$$\text{So } 2^{13} = 2^{n-1}$$

$$\therefore 13 = n - 1 \quad (\text{same bases})$$


$$\therefore n = 14$$

There are 16 384 newly-infected people on the 14<sup>th</sup> day.

For this geometric sequence, plotting the number of newly-infected people ( $T_n$ ) vs. the number of days ( $n$ ) results in the following graph:

Day ( $n$ )	No. of newly-infected people
1	2
2	4
3	8
4	16
5	32
6	64
$n$	$2 \times 2^{n-1}$





In this example we are only dealing with positive integers ( $n \in \{1; 2; 3; \dots\}, T_n \in \{1; 2; 3; \dots\}$ ) therefore the graph is not continuous and we do not join the points with a curve (the dotted line has been drawn to indicate the shape of an exponential graph).

### Geometric mean

The geometric mean between two numbers is the value that forms a geometric sequence together with the two numbers.

For example, the geometric mean between 5 and 20 is the number that has to be inserted between 5 and 20 to form the geometric sequence: 5;  $x$ ; 20

$$\begin{aligned} \text{Determine the constant ratio: } \frac{x}{5} &= \frac{20}{x} \\ \therefore x^2 &= 20 \times 5 \\ x^2 &= 100 \\ x &= \pm 10 \end{aligned}$$

---

**important:** Remember to include both the positive and negative square root. The geometric mean generates two possible geometric sequences:

$$5; 10; 20; \dots$$

$$5; -10; 20; \dots$$

In general, the geometric mean ( $x$ ) between two numbers  $a$  and  $b$  forms a geometric sequence with  $a$  and  $b$ :

For a geometric sequence:  $a; x; b$

Determine the constant ratio:  $\frac{x}{a} = \frac{b}{x}$

$$x^2 = ab$$

$$\therefore x = \pm\sqrt{ab}$$

## 3 SERIES

### 3.1 Series

It is often important and valuable to determine the sum of the terms of an arithmetic or geometric sequence. The sum of any sequence of numbers is called a series.

#### Finite series

We use the symbol  $S_n$  for the sum of the first  $n$  terms of a sequence  $\{T_1; T_2; T_3; \dots; T_n\}$

$$S_n = T_1 + T_2 + T_3 + \dots + T_n$$

If we sum only a finite number of terms, we get a finite series.

For example, consider the following sequence of numbers

$$1; 4; 9; 16; 25; 36; 49; \dots$$

We can calculate the sum of the first four terms:

$$S_4 = 1 + 4 + 9 + 16 = 30$$

This is an example of a finite series since we are only summing four terms.

#### Infinite series

If we sum infinitely many terms of a sequence, we get an infinite series:

$$S_\infty = T_1 + T_2 + T_3 + \dots$$

---

## 3.2 Sigma notation

Sigma notation is a very useful and compact notation for writing the sum of a given number of terms of a sequence.

A sum may be written out using the summation symbol  $\sum$  (Sigma), which is the capital letter “S” in the Greek alphabet. It indicates that you must sum the expression to the right of the summation symbol:

For example,

$$\sum_{n=1}^5 2n = 2 + 4 + 6 + 8 + 10 = 30$$

In general,

$$\sum_{i=m}^n T_i = T_m + T_{m+1} + \cdots + T_{n-1} + T_n$$

where

- $i$  is the index of the sum;
- $m$  is the lower bound (or start index), shown below the summation symbol;
- $n$  is the upper bound (or end index), shown above the summation symbol;
- $T_i$  is a term of a sequence;
- the number of terms in the series = end index – start index + 1.

The index  $i$  increases from  $m$  to  $n$  by steps of 1.

Note that this is also sometimes written as:

$$\sum_{i=m}^n a_i = a_m + a_{m+1} + \cdots + a_{n-1} + a_n$$

When we write out all the terms in a sum, it is referred to as the expanded form.

If we are summing from  $i = 1$  (which implies summing from the first term in a sequence), then we can use either  $S_n$  or  $\sum$  notation:

$$S_n = \sum_{i=1}^n a_i = a_1 + a_2 + \cdots + a_n \quad (n \text{ terms})$$

### WORKED EXAMPLE 4: SIGMA NOTATION

#### QUESTION

Expand the sequence and find the value of the series:

$$\sum_{n=1}^6 2^n$$

#### WORKED EXAMPLE 4: SIGMA NOTATION (continued)

##### SOLUTION

**Step 1: Expand the formula and write down the first six terms of the sequence**

$$\begin{aligned}\sum_{n=1}^6 2^n &= 2^1 + 2^2 + 2^3 + 2^4 + 2^5 + 2^6 \quad (6 \text{ terms}) \\ &= 2 + 4 + 8 + 16 + 32 + 64\end{aligned}$$

This is a geometric sequence 2; 4; 8; 16; 32; 64 with a constant ratio of 2 between consecutive terms.

**Step 2: Determine the sum of the first six terms of the sequence**

$$\begin{aligned}S_6 &= 2 + 4 + 8 + 16 + 32 + 64 \\ &= 126\end{aligned}$$

#### WORKED EXAMPLE 5: SIGMA NOTATION

##### QUESTION

Find the value of the series:

$$\sum_{n=3}^7 2an$$

##### SOLUTION

**Step 1: Expand the sequence and write down the five terms**

$$\begin{aligned}\sum_{n=3}^7 2an &= 2a(3) + 2a(4) + 2a(5) + 2a(6) + 2a(7) \quad (5 \text{ terms}) \\ &= 6a + 8a + 10a + 12a + 14a\end{aligned}$$

**Step 2: Determine the sum of the five terms of the sequence**

$$\begin{aligned}S_5 &= 6a + 8a + 10a + 12a + 14a \\ &= 50a\end{aligned}$$

#### WORKED EXAMPLE 6: SIGMA NOTATION

##### QUESTION

Write the following series in sigma notation:

$$31 + 24 + 17 + 10 + 3$$

##### SOLUTION

**Step 1: Consider the series and determine if it is an arithmetic or geometric series**

First test for an arithmetic series: is there a common difference?

### WORKED EXAMPLE 6: SIGMA NOTATION (continued)

We let:

$$T_1 = 31; \quad T_4 = 10;$$

$$T_2 = 24; \quad T_5 = 3;$$

$$T_3 = 17;$$

We calculate:

$$d = T_2 - T_1$$

$$= 24 - 31$$

$$= -7$$

$$d = T_3 - T_2$$

$$= 17 - 24$$

$$= -7$$

There is a common difference of  $-7$ , therefore this is an arithmetic series.

**Step 2: Determine the general formula of the series**

$$T_n = a + (n - 1)d$$

$$= 31 + (n - 1)(-7)$$

$$= 31 - 7n + 7$$

$$= -7n + 38$$

**Be careful:** brackets must be used when substituting  $d = -7$  into the general term. Otherwise the equation would be  $T_n = 31 + (n - 1) - 7$  which would be incorrect.

**Step 3: Determine the sum of the series and write in sigma notation**

$$31 + 24 + 17 + 10 + 3 = 85$$

$$\therefore \sum_{n=1}^5 (-7n + 38) = 85$$

### Rules for sigma notation

1. Given two sequences,  $a_i$  and  $b_i$ :

$$\sum_{i=1}^n (a_i + b_i) = \sum_{i=1}^n a_i + \sum_{i=1}^n b_i$$

---

2. For any constant  $c$  that is not dependent on the index  $i$ :

$$\begin{aligned}\sum_{i=1}^n (c \cdot a_i) &= c \cdot a_1 + c \cdot a_2 + c \cdot a_3 + \cdots + c \cdot a_n \\ &= c(a_1 + a_2 + a_3 + \cdots + a_n) \\ &= c \sum_{i=1}^n a_i\end{aligned}$$

3. Be accurate with the use of brackets:

Example 1:

$$\begin{aligned}\sum_{n=1}^3 (2n + 1) &= 3 + 5 + 7 \\ &= 15\end{aligned}$$

Example 2:

$$\begin{aligned}\sum_{n=1}^3 (2n) + 1 &= (2 + 4 + 6) + 1 \\ &= 13\end{aligned}$$

Note: the series in the second example has the general term  $T_n = 2n$  and the  $+1$  is added to the sum of the three terms. It is very important in sigma notation to use brackets correctly.

4.

$$\sum_{i=m}^n a_i$$

The values of  $i$ :

1. start at  $m$  ( $m$  is not always 1);
2. increase in steps of 1;
3. and end at  $n$ .

---

## 4 FINITE ARITHMETIC SERIES

### 4.1 Finite arithmetic series

An arithmetic sequence is a sequence of numbers, such that the difference between any term and the previous term is a constant number called the common difference ( $d$ ):

$$T_n = a + (n - 1)d$$

where

- $T_n$  is the  $n^{\text{th}}$  term of the sequence;
- $a$  is the first term;
- $d$  is the common difference.

When we sum a finite number of terms in an arithmetic sequence, we get a finite arithmetic series.

#### The sum of the first one hundred integers

A simple arithmetic sequence is when  $a = 1$  and  $d = 1$  which is the sequence of positive integers:

$$\begin{aligned}T_n &= a + (n - 1)d \\ &= 1 + (n - 1)(1) \\ &= n \\ \therefore \{T_n\} &= 1; 2; 3; 4; 5; \dots\end{aligned}$$

If we wish to sum this sequence from  $n = 1$  to any positive integer, for example 100, we would write

$$\sum_{n=1}^{100} n = 1 + 2 + 3 + \dots + 100$$

This gives the answer to the sum of the first 100 positive integers.

The mathematician, Karl Friedrich Gauss, discovered the following proof when he was only 8 years old. His teacher had decided to give his class a problem which would distract them for the entire day by asking them to add all the numbers from **1** to **100**. Young Karl quickly realised how to do this and shocked the teacher with the correct answer, **5050**. This is the method that he used:

- Write the numbers in ascending order.
- Write the numbers in descending order.

- Add the corresponding pairs of terms together.
- Simplify the equation by making  $S_n$  the subject of the equation.

$$\begin{aligned}
 S_{100} &= 1 + 2 + 3 + \dots + 98 + 99 + 100 \\
 + \quad S_{100} &= \underline{100 + 99 + 98 + \dots + 3 + 2 + 1} \\
 \therefore 2S_{100} &= 101 + 101 + 101 + \dots + 101 + 101 + 101 \\
 \therefore 2S_{100} &= 101 \times 100 \\
 &= 10\,100 \\
 \therefore S_{100} &= \frac{10100}{2} \\
 &= 5\,050
 \end{aligned}$$

## 4.2 General formula for a finite arithmetic series

If we sum an arithmetic sequence, it takes a long time to work it out term-by-term. We therefore derive the general formula for evaluating a finite arithmetic series. We start with the general formula for an arithmetic sequence of  $n$  terms and sum it from the first term ( $a$ ) to the last term in the sequence ( $l$ ) :

$$\begin{aligned}
 \sum_{n=1}^l T_n &= S_n \\
 S_n &= a + (a + d) + (a + 2d) + \dots + (l - 2d) + (l - d) + l \\
 + \quad S_n &= \underline{l + (l - d) + (l - 2d) + \dots + (a + 2d) + (a + d) + a} \\
 \therefore 2S_n &= (a + l) + (a + l) + (a + l) + \dots + (a + l) + (a + l) + (a + l) \\
 \therefore 2S_n &= n \times (a + l) \\
 \therefore S_n &= \frac{n}{2}(a + l)
 \end{aligned}$$

This general formula is useful if the last term in the series is known.

We substitute  $l = a + (n - 1)d$  into the above formula and simplify:

$$\begin{aligned}
 S_n &= \frac{n}{2}(a + [a + (n - 1)d]) \\
 \therefore S_n &= \frac{n}{2}[2a + (n - 1)d]
 \end{aligned}$$



---

The general formula for determining the sum of an arithmetic series is given by:

$$S_n = \frac{n}{2} [2a + (n - 1) d]$$

or

$$S_n = \frac{n}{2} (a + l)$$

For example, we can calculate the sum  $S_{20}$  for the arithmetic sequence  $T_n = 3 + 7(n - 1)$  by summing all the individual terms:

$$\begin{aligned} S_{20} &= \sum_{n=1}^{20} [3 + 7(n - 1)] \\ &= 3 + 10 + 17 + 24 + 31 + 38 + 45 + 52 \\ &\quad + 59 + 66 + 73 + 80 + 87 + 94 + 101 \\ &\quad + 108 + 115 + 122 + 129 + 136 \\ &= 1\,390 \end{aligned}$$

or, more sensibly, we could use the general formula for determining an arithmetic series by substituting  $a = 3$ ,  $d = 7$  and  $n = 20$ :

$$\begin{aligned} S_n &= \frac{n}{2} (2a + (n - 1)d) \\ S_{20} &= \frac{20}{2} [2(3) + 7(20 - 1)] \\ &= 1\,390 \end{aligned}$$

This example demonstrates how useful the general formula for determining an arithmetic series is, especially when the series has a large number of terms.

### WORKED EXAMPLE 7: GENERAL FORMULA FOR THE SUM OF AN ARITHMETIC SERIES

**QUESTION** Find the sum of the first 30 terms of an arithmetic series with  $T_n = 7n - 5$  by using the formula.

**SOLUTION**

**Step 1: Use the general formula to generate terms of the sequence and write down the known variables**

$$\begin{aligned}T_n &= 7n - 5 \\ \therefore T_1 &= 7(1) - 5 \\ &= 2 \\ T_2 &= 7(2) - 5 \\ &= 9 \\ T_3 &= 7(3) - 5 \\ &= 16\end{aligned}$$

This gives the sequence: 2; 9; 16 ...

$$a = 2; \quad d = 7; \quad n = 30$$

### WORKED EXAMPLE 7: GENERAL FORMULA FOR THE SUM OF AN ARITHMETIC SERIES (continued)

**Step 2: Write down the general formula and substitute the known values**

$$\begin{aligned}S_n &= \frac{n}{2}[2a + (n - 1)d] \\ S_{30} &= \frac{30}{2}[2(2) + (30 - 1)(7)] \\ &= 15(4 + 203) \\ &= 15(207) \\ &= 3\,105\end{aligned}$$

**Step 3: Write the final answer**

$$S_{30} = 3\,105$$

### WORKED EXAMPLE 8: SUM OF AN ARITHMETIC SERIES IF FIRST AND LAST TERMS ARE KNOWN

#### QUESTION

Find the sum of the series  $-5 - 3 - 1 + \dots + 123$ .

#### SOLUTION

**Step 1: Identify the type of series and write down the known variables**

$$\begin{aligned}d &= T_2 - T_1 \\ &= -3 - (-5) \\ &= 2\end{aligned}$$

$$\begin{aligned}d &= T_3 - T_2 \\ &= -1 - (-3) \\ &= 2\end{aligned}$$

$$a = -5; \quad d = 2; \quad l = 123$$

**Step 2: Determine the value of  $n$**

$$\begin{aligned}T_n &= a + (n - 1)d \\ \therefore 123 &= -5 + (n - 1)(2) \\ &= -5 + 2n - 2 \\ \therefore 130 &= 2n \\ \therefore n &= 65\end{aligned}$$

**Step 3: Use the general formula to find the sum of the series**

$$\begin{aligned}S_n &= \frac{n}{2}(a + l) \\ S_{65} &= \frac{65}{2}(-5 + 123) \\ &= \frac{65}{2}(118) \\ &= 3\,835\end{aligned}$$

**Step 4: Write the final answer**

$$S_{65} = 3\,835$$

### WORKED EXAMPLE 9: FINDING $n$ GIVEN THE SUM OF AN ARITHMETIC SEQUENCE

#### QUESTION

Given an arithmetic sequence with  $T_2 = 7$  and  $d = 3$ , determine how many terms must be added together to give a sum of 2 146.

#### SOLUTION

**Step 1: Write down the known variables**

$$d = T_2 - T_1$$

$$\therefore 3 = 7 - a$$

$$\therefore a = 4$$

$$a = 4; \quad d = 3; \quad S_n = 2\,146$$

**Step 2: Use the general formula to determine the value of  $n$**

$$S_n = \frac{n}{2}(2a + (n-1)d)$$

$$2146 = \frac{n}{2}(2(4) + (n-1)(3))$$

$$4292 = n(8 + 3n - 3)$$

$$\therefore 0 = 3n^2 + 5n - 4292$$

$$= (3n + 116)(n - 37)$$

$$\therefore n = -\frac{116}{3} \text{ or } n = 37$$

but  $n$  must be a positive integer, therefore  $n = 37$ .

We could have solved for  $n$  using the quadratic formula but factorising by inspection is usually the quickest method.

**Step 3: Write the final answer**

$$S_{37} = 2\,146$$

### WORKED EXAMPLE 10: FINDING $n$ GIVEN THE SUM OF AN ARITHMETIC SERIES

#### QUESTION

The sum of the second and third terms of an arithmetic sequence is equal to zero and the sum of the first 36 terms of the series is equal to 1 152. Find the first three terms in the series.

#### SOLUTION

##### Step 1: Write down the given information

$$S_n = \frac{n}{2}(2a + (n - 1)d)$$

$$S_{36} = \frac{36}{2}(2a + (36 - 1)d)$$

$$1152 = 18(2a + 35d)$$

$$\therefore 64 = 2a + 35d \dots\dots (2)$$

##### Step 2: Solve the two equations simultaneously

$$2a + 3d = 0 \dots\dots (1)$$

$$2a + 35d = 64 \dots\dots (2)$$

$$\text{Eqn (2) - (1) : } 32d = 64$$

$$\therefore d = 2$$

$$\text{And } 2a + 3(2) = 0$$

$$2a = -6$$

$$\therefore a = -3$$

##### Step 3: Write the final answer The first three terms of the series are:

$$T_1 = a = -3$$

$$T_2 = a + d = -3 + 2 = -1$$

$$T_3 = a + 2d = -3 + 2(2) = 1$$

$$-3 - 1 + 1$$

---

### Calculating the value of a term given the sum of $n$ terms:

If the first term in a series is  $T_1$ , then  $S_1 = T_1$ .

We also know the sum of the first two terms  $S_2 = T_1 + T_2$ , which we rearrange to make  $T_2$  the subject of the equation:

$$T_2 = S_2 - T_1$$

Substitute  $S_1 = T_1$

$$\therefore T_2 = S_2 - S_1$$

Similarly, we could determine the third and fourth term in a series:

$$T_3 = S_3 - S_2$$

$$\text{And } T_4 = S_4 - S_3$$

$$T_n = S_n - S_{n-1}, \text{ for } n \in \{2; 3; 4; \dots\} \text{ and } T_1 = S_1.$$

## 5 FINITE GEOMETRIC SERIES

### 5.1 Finite geometric series

When we sum a known number of terms in a geometric sequence, we get a finite geometric series. We generate a geometric sequence using the general form:

$$T_n = a \cdot r^{n-1}$$

where

- $n$  is the position of the sequence;
- $T_n$  is the  $n^{\text{th}}$  term of the sequence;
- $a$  is the first term;
- $r$  is the constant ratio.

---

## 5.2 General formula for a finite geometric series

$$S_n = a + ar + ar^2 + \dots + ar^{n-2} + ar^{n-1} \dots (1)$$

$$r \times S_n = ar + ar^2 + \dots + ar^{n-2} + ar^{n-1} + ar^n \dots (2)$$

Subtract eqn. (2) from eqn. (1)

$$\therefore S_n - rS_n = a + 0 + 0 + \dots - ar^n$$

$$S_n - rS_n = a - ar^n$$

$$S_n(1 - r) = a(1 - r^n)$$

$$\therefore S_n = \frac{a(1 - r^n)}{1 - r} \quad (\text{where } r \neq 1)$$

The general formula for determining the sum of a geometric series is given by:

$$S_n = \frac{a(1-r^n)}{1-r} \quad \text{where } r \neq 1$$

This formula is easier to use when  $r < 1$

**Alternative formula:**

$$S_n = a + ar + ar^2 + \dots + ar^{n-2} + ar^{n-1} \dots (1)$$

$$r \times S_n = ar + ar^2 + \dots + ar^{n-2} + ar^{n-1} + ar^n \dots (2)$$

Subtract eqn. (1) from eqn. (2)

$$\therefore rS_n - S_n = ar^n - a$$

$$S_n(r - 1) = a(r^n - 1)$$

$$\therefore S_n = \frac{a(r^n - 1)}{r - 1} \quad (\text{where } r \neq 1)$$

The general formula for determining the sum of a geometric series is given by:

$$S_n = \frac{a(r^n-1)}{r-1} \quad \text{where } r \neq 1$$

This formula is easier to use when  $r > 1$ .

### Example : Number Formats and Conventions

#### QUESTION

Calculate:

$$\sum_{k=1}^6 32 \left(\frac{1}{2}\right)^{k-1}$$

#### SOLUTION

**Step 1: Write down the first three terms of the series**

$$k = 1; \quad T_1 = 32 \left(\frac{1}{2}\right)^0 = 32$$

$$k = 2; \quad T_2 = 32 \left(\frac{1}{2}\right)^{2-1} = 16$$

$$k = 3; \quad T_3 = 32 \left(\frac{1}{2}\right)^{3-1} = 8$$

We have generated the series  $32 + 16 + 8 + \dots$

**Step 2: Determine the values of a and r**

$$a = T_1 = 32$$

$$r = \frac{T_2}{T_1} = \frac{T_3}{T_2} = \frac{1}{2}$$

**Step 3: Use the general formula to find the sum of the series**

$$S_n = \frac{a(1 - r^n)}{1 - r}$$

$$S_6 = \frac{32(1 - (\frac{1}{2})^6)}{1 - \frac{1}{2}}$$

$$= \frac{32(1 - \frac{1}{64})}{\frac{1}{2}}$$

$$= 2 \times 32 \left(\frac{63}{64}\right)$$

$$= 64 \left(\frac{63}{64}\right)$$

$$= 63$$

**Step 4: Write the final answer**

$$\sum_{k=1}^6 32 \left(\frac{1}{2}\right)^{k-1} = 63$$



## WORKED EXAMPLE 12: SUM OF A GEOMETRIC SERIES

### QUESTION

Given a geometric series with  $T_1 = -4$  and  $T_4 = 32$ . Determine the values of  $r$  and  $n$  if  $S_n = 84$ .

### SOLUTION

**Step 1: Determine the values of  $a$  and  $r$**

$$\begin{aligned}a &= T_1 = -4 \\T_4 &= ar^3 = 32 \\ \therefore -4r^3 &= 32 \\ r^3 &= -8 \\ \therefore r &= -2\end{aligned}$$

Therefore the geometric series is  $-4 + 8 - 16 + 32 \dots$ . Notice that the signs of the terms alternate because  $r < 0$ .

We write the general term for this series as  $T_n = -4(-2)^{n-1}$

**Step 2: Use the general formula for the sum of a geometric series to determine the value of  $n$**

$$\begin{aligned}S_n &= \frac{a(1 - r^n)}{1 - r} \\ \therefore 84 &= \frac{-4(1 - (-2)^n)}{1 - (-2)} \\ 84 &= \frac{-4(1 - (-2)^n)}{3} \\ -\frac{3}{4} \times 84 &= 1 - (-2)^n \\ -63 &= 1 - (-2)^n \\ (-2)^n &= 64 \\ (-2)^n &= (-2)^6 \\ \therefore n &= 6\end{aligned}$$

**Step 3: Write the final answer**

$r = -2$  and  $n = 6$

### WORKED EXAMPLE 13: SUM OF A GEOMETRIC SERIES

#### QUESTION

Use the general formula for the sum of a geometric series to determine  $k$  if

$$\sum_{n=1}^8 k \left(\frac{1}{2}\right)^n = \frac{255}{64}$$

#### SOLUTION

**Step 1: Write down the first three terms of the series**

$$n = 1; \quad T_1 = k \left(\frac{1}{2}\right)^1 = \frac{1}{2}k$$

$$n = 2; \quad T_2 = k \left(\frac{1}{2}\right)^2 = \frac{1}{4}k$$

$$n = 3; \quad T_3 = k \left(\frac{1}{2}\right)^3 = \frac{1}{8}k$$

We have generated the series  $\frac{1}{2}k + \frac{1}{4}k + \frac{1}{8}k + \dots$

We can take out the common factor  $k$  and write the series as:  $k \left(\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots\right)$

$$\therefore k \sum_{n=1}^8 \left(\frac{1}{2}\right)^n = \frac{255}{64}$$

**Step 2: Determine the values of  $a$  and  $r$**

$$a = T_1 = \frac{1}{2}$$

$$r = \frac{T_2}{T_1} = \frac{T_3}{T_2} = \frac{1}{2}$$

**Step 3: Calculate the sum of the first eight terms of the geometric series**

$$\therefore S_n = \frac{a(1 - r^n)}{1 - r}$$

$$S_8 = \frac{\frac{1}{2}(1 - (\frac{1}{2})^8)}{1 - \frac{1}{2}}$$

$$= \frac{\frac{1}{2}(1 - (\frac{1}{2})^8)}{\frac{1}{2}}$$

$$= 1 - \frac{1}{256}$$

$$= \frac{255}{256}$$

$$\therefore \sum_{n=1}^8 \left(\frac{1}{2}\right)^n = \frac{255}{256}$$

### WORKED EXAMPLE 13: SUM OF A GEOMETRIC SERIES (continued)

So then we can write:

$$\begin{aligned}k \sum_{n=1}^8 \left(\frac{1}{2}\right)^n &= \frac{255}{64} \\k \left(\frac{255}{256}\right) &= \frac{255}{64} \\ \therefore k &= \frac{255}{64} \times \frac{256}{255} \\ &= \frac{256}{64} \\ &= 4\end{aligned}$$

**Step 4: Write the final answer**

$$k = 4$$

## 6 INFINITE SERIES

### 6.1 Infinite series

So far we have been working only with finite sums, meaning that whenever we determined the sum of a series, we only considered the sum of the first  $n$  terms. We now consider what happens when we add an infinite number of terms together. Surely if we sum infinitely many numbers, no matter how small they are, the answer goes to infinity? In some cases the answer does indeed go to infinity (like when we sum all the positive integers), but surprisingly there are some cases where the answer is a finite real number.

#### INVESTIGATION

##### Sum of an infinite series

- Cut a piece of string 1m in length.
- Now cut the piece of string in half and place one half on the desk.
- Cut the other half in half again and put one of the pieces on the desk.
- Repeat this process until the piece of string is too short to cut easily.

### INVESTIGATION (continued)

- Draw a diagram to illustrate the sequence of lengths of the pieces of string.
- Can this sequence be expressed mathematically? Hint: express the shorter lengths of string as a fraction of the original length of string.
- What is the sum of the lengths of all the pieces of string?
- Predict what would happen if these steps could be repeated infinitely many times.
- Will the sum of the lengths of string ever be greater than 1?
- What can you conclude?

### WORKED EXAMPLE 14: SUM TO INFINITY

#### QUESTION

Complete the table below for the geometric series  $T_n = \left(\frac{1}{2}\right)^n$  and answer the questions that follow:

	Terms	$S_n$	$1 - S_n$
$T_1$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$
$T_1 + T_2$			
$T_1 + T_2 + T_3$			
$T_1 + T_2 + T_3 + T_4$			

1. As more and more terms are added, what happens to the value of  $S_n$ ?
2. As more and more terms are added, what happens to the value of  $1 - S_n$ ?
3. Predict the maximum value of  $S_n$  for the sum of infinitely many terms in the series.

#### SOLUTION

##### Step 1: Complete the table

	Terms	$S_n$	$1 - S_n$
$T_1$	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$
$T_1 + T_2$	$\frac{1}{2} + \frac{1}{4}$	$\frac{3}{4}$	$\frac{1}{4}$
$T_1 + T_2 + T_3$	$\frac{1}{2} + \frac{1}{4} + \frac{1}{8}$	$\frac{7}{8}$	$\frac{1}{8}$
$T_1 + T_2 + T_3 + T_4$	$\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \frac{1}{16}$	$\frac{15}{16}$	$\frac{1}{16}$

### WORKED EXAMPLE 14: SUM TO INFINITY (continued)

#### Step 2: Consider the value of $S_n$ and $1 - S_n$

As more terms in the series are added together, the value of  $S_n$  increases:

$$\frac{1}{2} < \frac{3}{4} < \frac{7}{8} < \dots$$

However, by considering  $1 - S_n$  we notice that the amount by which  $S_n$  increases gets smaller and smaller as more terms are added:

$$\frac{1}{2} > \frac{1}{4} > \frac{1}{8} > \dots$$

We can therefore conclude that the value of  $S_n$  is approaching a maximum value of **1**; it is converging to **1**.

#### Step 3: Write conclusion mathematically

We can conclude that the sum of the series

$$\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots$$

gets closer to 1.  $S_n \rightarrow 1$  as the number of terms approaches infinity ( $n \rightarrow \infty$ ), therefore the series converges.

$$\sum_{i=1}^{\infty} \left(\frac{1}{2}\right)^i = 1$$

We express the sum of an infinite number of terms of a series as

$$S_{\infty} = \sum_{i=1}^{\infty} T_i$$

### Convergence and divergence

If the sum of a series gets closer and closer to a certain value as we increase the number of terms in the sum, we say that the series converges. In other words, there is a limit to the sum of a converging series. If a series does not converge, we say that it diverges. The sum of an infinite series usually tends to infinity, but there are some special cases where it does not.

Note the following:

- An arithmetic series never converges: as  $n$  tends to infinity, the series will always tend to positive or negative infinity.
- Some geometric series converge (have a limit) and some diverge (as  $n$  tends to infinity, the series does not tend to any limit or it tends to infinity).

---

## 6.2 Infinite geometric series

There is a simple test for determining whether a geometric series converges or diverges; if  $-1 < r < 1$  then the infinite series will converge. If  $r$  lies outside this interval, then the infinite series will diverge.

### Test for convergence:

- If  $-1 < r < 1$ , then the infinite geometric series converges.
- If  $r < -1$  or  $r > 1$ , then the infinite geometric series diverges.

We derive the formula for calculating the value to which a geometric series converges as follows:

$$S_n = \sum_{i=1}^n ar^{i-1} = \frac{a(1-r^n)}{1-r}$$

Now consider the behaviour of  $r^n$  for  $-1 < r < 1$  as  $n$  becomes larger.

Let  $r = \frac{1}{2}$ :

$$n = 1 : r^n = r^1 = \left(\frac{1}{2}\right)^1 = \frac{1}{2}$$

$$n = 2 : r^n = r^2 = \left(\frac{1}{2}\right)^2 = \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{4} < \frac{1}{2}$$

$$n = 3 : r^n = r^3 = \left(\frac{1}{2}\right)^3 = \frac{1}{2} \cdot \frac{1}{2} \cdot \frac{1}{2} = \frac{1}{8} < \frac{1}{4}$$

Since  $r$  is in the range  $-1 < r < 1$  we see that  $r^n$  gets closer to 0 as  $n$  gets larger. Therefore  $(1 - r^n)$  gets closer to 1.

Therefore,

$$S_n = \frac{a(1-r^n)}{1-r}$$

If  $-1 < r < 1$ , then  $r^n \rightarrow 0$  as  $n \rightarrow \infty$

$$\begin{aligned} \therefore S_\infty &= \frac{a(1-0)}{1-r} \\ &= \frac{a}{1-r} \end{aligned}$$

The sum of an infinite geometric series is given by the formula

$$\therefore S_\infty = \sum_{i=1}^{\infty} ar^{i-1} = \frac{a}{1-r} \quad (-1 < r < 1)$$

where

- $a$  is the first term of the series;
- $r$  is the constant ratio.

Alternative notation:

$$\underbrace{S_n}_{n \rightarrow \infty} \rightarrow \frac{a}{1-r} \quad \text{if } -1 < r < 1$$

In words: as the number of terms ( $n$ ) tends to infinity, the sum of a converging geometric series ( $S_n$ ) tends to the value  $\frac{a}{1-r}$ .

### WORKED EXAMPLE 15: SUM TO INFINITY OF A GEOMETRIC SERIES

#### QUESTION

Given the series  $18 + 6 + 2 + \dots$ . Find the sum to infinity if it exists.

#### SOLUTION

##### Step 1: Determine the value of $r$

We need to know the value of  $r$  to determine whether the series converges or diverges.

$$\begin{aligned}\frac{T_2}{T_1} &= \frac{6}{18} \\ &= \frac{1}{3} \\ \frac{T_3}{T_2} &= \frac{2}{6} \\ &= \frac{1}{3} \\ \therefore r &= \frac{1}{3}\end{aligned}$$

Since,  $-1 < r < 1$  we can conclude that this is a convergent geometric series.

##### Step 2: Determine the sum to infinity

Write down the formula for the sum to infinity and substitute the known values:

$$a = 18; \quad r = \frac{1}{3}$$

$$\begin{aligned}S_\infty &= \frac{a}{1-r} \\ &= \frac{18}{1-\frac{1}{3}} \\ &= \frac{18}{\frac{2}{3}} \\ &= 18 \times \frac{3}{2} \\ &= 27\end{aligned}$$

As  $n$  tends to infinity, the sum of this series tends to 27; no matter how many terms are added together, the value of the sum will never be greater than 27.

## WORKED EXAMPLE 16: USING THE SUM TO INFINITY TO CONVERT RECURRING DECIMALS TO FRACTIONS

### QUESTION

Use two different methods to convert the recurring decimal  $0,\dot{5}$  to a proper fraction.

### SOLUTION

#### Step 1: Convert the recurring decimal to a fraction using equations

$$\text{Let } x = 0,\dot{5}$$

$$\therefore x = 0,555\dots\dots(1)$$

$$10x = 5,55\dots\dots(2)$$

$$(2) - (1): \quad 9x = 5$$

$$\therefore x = \frac{5}{9}$$

#### Step 2: Convert the recurring decimal to a fraction using the sum to infinity

$$0,\dot{5} = 0,5 + 0,05 + 0,005 + \dots$$

$$\text{or } 0,\dot{5} = \frac{5}{10} + \frac{5}{100} + \frac{5}{1000} + \dots$$

This is a geometric series with  $r = 0,1 = \frac{1}{10}$  and since  $-1 < r < 1$ , we can conclude that the series is convergent.

$$\begin{aligned} S_{\infty} &= \frac{a}{1-r} \\ &= \frac{\frac{5}{10}}{1 - \frac{1}{10}} \\ &= \frac{\frac{5}{10}}{\frac{9}{10}} \\ &= \frac{5}{9} \end{aligned}$$



### WORKED EXAMPLE 17: USING THE SUM TO INFINITY TO CONVERT RECURRING DECIMALS TO FRACTIONS

#### QUESTION

Determine the possible values of  $a$  and  $r$  if

$$\sum_{n=1}^{\infty} ar^{n-1} = 5$$

#### SOLUTION

**Step 1: Write down the sum to infinity formula and substitute known values**

$$S_{\infty} = \frac{a}{1-r}$$

$$\therefore 5 = \frac{a}{1-r}$$

$$a = 5(1-r)$$

$$\therefore a = 5 - 5r$$

$$\text{And } 5r = 5 - a$$

$$\therefore r = \frac{5-a}{5}$$

**Step 2: Apply the condition for convergence to determine possible values of  $a$**

For a series to converge:  $-1 < r < 1$

$$-1 < r < 1$$

$$-1 < \frac{5-a}{5} < 1$$

$$-5 < 5-a < 5$$

$$-10 < -a < 0$$

$$0 < a < 10$$

**Step 3: Write the final answer**

For the series to converge,  $0 < a < 10$  and  $-1 < r < 1$ .

## 7 SUMMARY

### Arithmetic sequence

- common difference ( $d$ ) between any two consecutive terms:  $d = T_n - T_{n-1}$
- general form:  $a + (a + d) + (a + 2d) + \dots$
- general formula:  $T_n = a + (n - 1)d$
- graph of the sequence lies on a straight line

---

### Quadratic sequence

- common second difference between any two consecutive terms
- general formula:  $T_n = an^2 + bn + c$
- graph of the sequence lies on a parabola

### Geometric sequence

- constant ratio ( $r$ ) between any two consecutive terms:  $r = \frac{T_n}{T_{n-1}}$
- general form:  $a + ar + ar^2 + \dots$
- general formula:  $T_n = ar^{n-1}$
- graph of the sequence lies on an exponential curve

### Sigma notation

$$\sum_{k=1}^n T_k$$

Sigma notation is used to indicate the sum of the terms given by  $T_k$  starting from  $k = 1$  and ending at  $k = n$ .

### Series

- the sum of certain numbers of terms in a sequence

- arithmetic series:

$$- S_n = \frac{n}{2}[a + l]$$

$$- S_n = \frac{n}{2}[2a + (n - 1)d]$$

- geometric series:

$$- S_n = \frac{a(1-r^n)}{1-r}$$

$$- S_n = \frac{a(r^n-1)}{r-1}$$

### Sum to infinity

A convergent geometric series, with  $-1 < r < 1$  tends to a certain fixed number as the number of terms in the sum tends to infinity.

$$S_\infty = \sum_{n=1}^{\infty} T_n = \frac{a}{1-r}$$

---

## 8 EXERCISES

### 8.1 Exercise 1

1. Find the common difference and write down the next three terms of the sequence.

1.1  $-5; -3; -1; 1; 3; \dots$

1.2  $-1; 10; 21; 32; 43; 54; \dots$

2. Given the sequence:  $2; 6; 10; 14; 18; 22; \dots$

2.1 Show that the above sequence is an Arithmetic sequence.

2.2 Write down the next three terms in the sequence.

2.3 Determine the general term of the sequence.

2.4 Which term has a value of 86?

2.5 Prove that 236 is not a term in this sequence.

3. Given the sequence:  $-1; -4; -7; -10; -13; -16; \dots$

3.1 Show that the above sequence is an Arithmetic sequence.

3.2 Write down the next three terms in the sequence.

3.3 Calculate the value of the  $20^{\text{th}}$  term.

3.4 Which term has a value of  $-94$ ?

3.5 Prove that  $-71$  is not a term in this sequence.

4. Given the sequence:  $-2; -\frac{3}{2}; -1; -\frac{1}{2}; 0; \frac{1}{2}; 1; \dots$

4.1 Show that the above sequence is an Arithmetic sequence.

4.2 Write down the next three terms in the sequence.

4.3 Determine  $T_{100}$ .

4.4 Which term has a value of  $\frac{63}{2}$ ?

5. Given the sequence  $7; 5, 5; 4; 2, 5; \dots$


5.1 Find the next term in the sequence.

5.2 Determine the general term of the sequence.

5.3 Which term has a value of  $-23$ ?

- 
6. Given the sequence: 2; 6; 10; 14; ...
- 6.1 Is this an arithmetic sequence? Justify your answer by calculation.
  - 6.2 Calculate  $T_{55}$ .
  - 6.3 which term has the value of 322?
  - 6.4 Determine by calculation whether or not 1 204 is a term of this sequence.
7. An arithmetic sequence has the general term  $T_n = -2n + 7$ .
- 7.1 Calculate the second, third and tenth terms of the sequence.
  - 7.2 Draw a diagram of the sequence for  $0 < n \leq 10$ .
8. Given the arithmetic sequence: 2;  $x$ ; 12
- 8.1 Determine the value if  $x$ .
  - 8.2 Determine the position and value of the first term in the sequence greater than 100.
9. Given the arithmetic sequence:  $3x - 4$ ;  $4x - 2$ ;  $7x - 6$ . Determine the value of  $x$ .
10. Given the arithmetic sequence:  $a - 3b$ ;  $a - b$ ;  $a + b$ ;  $a + 3b$ ; ...
- 10.1 Prove that the above sequence is an arithmetic sequence.
  - 10.2 Write down the next term in the sequence.
  - 10.3 Determine the value of  $a$  and  $b$ , if  $T_1 = 2$  and  $T_4 = 14$ .
  - 10.4 Determine the value of  $T_{1\ 099}$ ?
11. The terms  $p$ ;  $(2p + 2)$ ;  $(5p + 3)$  form an arithmetic sequence. Find  $p$  and the  $15^{th}$  term of the sequence.
12. What are the important characteristics of an arithmetic sequence?
13. You are given the first four terms of an arithmetic sequence.  
Describe the method you would use to find the formula for the  $n^{th}$  term of the sequence.
14. A single square is made from 4 matchsticks. To make two squares in a row takes 7 matchsticks, while three squares in a row takes 10 matchsticks
- 14.1 Write down the first four terms of the sequence.
  - 14.2 What is the common difference?
  - 14.3 Determine the formula for the general term.
  - 14.4 How many matchsticks are in a row of 25 squares?
  - 14.5 if there are 109 matchsticks, calculate the number of squares in the row.

15. A pattern of equilateral triangles decorates the border of a girl's skirt. Each triangle is made by three stitches, each having a length of 1 cm.



- 15.1 Complete the table:

Figure no.	1	2	3	$q$	$r$	$n$
No. of stitches	3	5	$p$	15	71	$s$

- 15.2 The border of the skirt is 2 m in length. If the entire length of the border is decorated with the triangular pattern, how many stitches will there be?
16. In an arithmetic sequence the seventh term is 30 and the fourth term is 18. Determine the value of the first three terms in the sequence.
17. The first term of an arithmetic sequence is  $-\frac{1}{2}$  and  $T_{22} = 10$ . Find  $T_n$ .
18. In a arithmetic sequence the seventh term is 9 more than the fourth term. The sixth term is 5. Determine  $T_1$  and the constant difference.
19. The  $k^{th}$  term of an arithmetic sequence is  $m$  and the  $m^{th}$  term is equal to  $k$ . Find the common difference of the sequence.
20. Determine the number of terms in the following sequence:  $53 + 49 + 45 + \dots + 13$
21. The arithmetic mean of  $3a - 2$  and  $x$  is  $4a - 4$ . Determine the value of  $x$  in terms of  $a$ .
22. Insert seven arithmetic means between the terms  $(3s - t)$  and  $(-13s + 7t)$ .

## 8.2 Exercise 2

1. Determine whether each of the following sequences is:

- a linear sequence;
- a quadratic sequence;
- or neither

---

1.1 8; 17; 32; 53; 80; ...

1.2  $3p^2$ ;  $6p^2$ ;  $9p^2$ ;  $12p^2$ ;  $15p^2$ ; ...

1.3 1; 2,5; 5; 8, 5; 13; ...

1.4 2; 6; 10; 14; 18; ...

1.5 5; 19; 41; 71; 109; ...

1.6 3; 9; 16; 21; 27; ...

1.7  $2k$ ;  $8k$ ;  $18k$ ;  $32k$ ;  $50k$ ; ...

1.8  $2\frac{1}{2}$ ; 6;  $10\frac{1}{2}$ ; 16;  $22\frac{1}{2}$ ; ...

2. A quadratic pattern is given by  $T_n = n^2 + bn + c$ . Find the values of  $b$  and  $c$  if the sequence starts with the following terms:

-1; 2; 7; 14; ...

3.  $a^2$ ;  $-a^2$ ;  $-3a^2$ ;  $-5a^2$ ; ... are the first 4 terms of a sequence.

3.1 Is the sequence linear or quadratic?

3.2 What is the next term in the sequence?

3.3 Calculate  $T_{100}$

4. Given  $T_n = n^2 + bn + c$ , determine the values of  $b$  and  $c$  if the sequence starts with the terms:

2; 7; 14; 23; ...

5. The first term of a quadratic sequence is 4, the third term is 34 and the common second difference is 10. Determine the first six terms in the sequence.

6. A quadratic sequence has a second term equal to 1, a third term equal to -6 and a fourth term equal to -14.

6.1 Determine the second difference for this sequence.

6.2 Hence, or otherwise, calculate the first term of the pattern.

### 8.3 Exercise 3

1. Determine the constant ratios for the following geometric sequences and write down the next three terms in each sequence:

1.1 5; 10; 20; ...

1.2  $\frac{1}{2}$ ;  $\frac{1}{4}$ ;  $\frac{1}{8}$ ; ...

1.3 7; 0, 7; 0, 07; ...

1.4  $p$ ;  $3p^2$ ;  $9p^3$ ; ...

1.5  $-3; 30; -300; \dots$

2. Determine the general formula for the  $n^{\text{th}}$  term of each of the following geometric sequences:

2.1  $5; 10; 20; \dots$

2.2  $\frac{1}{2}; \frac{1}{4}; \frac{1}{8}; \dots$

2.3  $7; 0, 7; 0, 07; \dots$

2.4  $p; 3p^2; 9p^3; \dots$

2.5  $-3; 30; -300; \dots$

3. The  $n^{\text{th}}$  term of a sequence is given by the formula  $T_n = 6\left(\frac{1}{3}\right)^{n-1}$ .

3.1 Write down the first three terms of the sequence.

3.2 What type of sequence is this?

4. Consider the following terms:

$(k - 4); (k + 1); m; 5k$

The first three terms form an arithmetic sequence and the last three terms form a geometric sequence.

Determine the values of  $k$  and  $m$  if both are positive integers.


5. Given a geometric sequence with second term  $\frac{1}{2}$  and the ninth term 64.

5.1 Determine the value of  $r$ .

5.2 Find the value of  $a$ .

5.3 Determine the general formula of the sequence.

6. The diagram shows four sets of values of consecutive terms of a geometric sequence with the general formula  $T_n = ar^{n-1}$



- 
- 6.1 Determine  $a$  and  $r$ .
- 6.2 Find  $x$  and  $y$ .
- 6.3 Find the fifth term in the sequence.
7. Write down the next two terms for the following sequence:  $1; \sin \theta; 1 - \cos^2 \theta; \dots$
8.  $5; x; y$  is an arithmetic sequence and  $x; y; 81$  is a geometric sequence. All terms in the sequences are integers. Calculate the values of  $x$  and  $y$ .
9. The two numbers  $2x^2y^2$  and  $8x^4$  are given.
- 9.1 Write down the geometric mean between the two numbers in terms of  $x$  and  $y$ .
- 9.2 Determine the constant ratio of the resulting sequence.
10. Insert three geometric means between  $-1$  and  $-\frac{1}{81}$  and give all possible answers.

## 8.4 Exercise 4

1. Determine the value of the following:

1.1  $\sum_{k=1}^4 2$

1.2  $\sum_{i=-1}^3 i$

1.3  $\sum_{n=2}^5 (3n - 2)$

1.4  $\sum_{t=4}^7 3t^3$

1.5  $\sum_{k=0}^5 \left( \sin \left( \frac{\pi k}{2} \right) \right)$



---

2. Determine the series:

2.1  $\sum_{k=1}^6 0^k$

2.2  $\sum_{n=1}^4 8$

2.3  $\sum_{k=1}^5 (ak)$

2.4  $\sum_{x=0}^4 (x^2 - 3x + 1)$

3. Calculate the value of  $a$ .

3.1  $\sum_{k=1}^3 (a \cdot 2^{k-1}) = 28$

3.2  $\sum_{j=1}^4 (2^{-j}) = a$

3.3  $\sum_{n=0}^4 (-2)^k = a$

4. Write the following in sigma notation:

4.1  $\frac{1}{9} + \frac{1}{3} + 1 + 3$

4.2  $-\frac{1}{22} + \frac{1}{23} - \frac{1}{24} + \frac{1}{25} - \frac{1}{26}$

5. Write the sum of the first 25 terms of the series below in sigma notation:  $11 + 4 - 3 - 10 \dots$

6. Write the sum of the first 1 000 natural, odd numbers in sigma notation.

7. Evaluate in terms of  $n$ :

7.1  $\sum_{k=1}^n T_k$  if given a geometric sequence of 25; 5; 1 ...

7.2  $\sum_{x=1}^n Tr$  if given a geometric sequence of:  $-\frac{1}{4}$ ;  $\frac{1}{6}$ ;  $-\frac{1}{9} \dots$

## 8.5 Exercise 5

1. Calculate the sum of the first 15 terms of the series:

1.1  $3 + 7 + 11 + 15 + \dots$

1.2  $2 + 7 + 12 + 17 + \dots$

1.3  $-25 - 22 - 19 - 16 + \dots$

2. Calculate the sum of the arithmetic series  $4 + 7 + 10 + \dots + 901$

3. Calculate the sum of the arithmetic series  $7 + 10 + 13 + 16 \dots + 574$

4. Evaluate without using a calculator:

$$\frac{4+8+12+\dots+100}{3+10+17+\dots+101}$$

---

5. Consider the following series:

$$-19 - 14 - 9 - 4 \dots + 976$$

5.1 How many terms are in the series?

5.2 Calculate the sum of the series.

6. The sum to  $n$  terms of an arithmetic series is  $S_n = \frac{n}{2}(7n + 15)$

6.1 How many terms of the series must be added to give a sum of 425?

6.2 Determine the sixth term of the series.

7. Determine the value of the following:

7.1  $\sum_{w=0}^8 (7w + 8)$

7.2  $\sum_{j=1}^8 7j + 8$

8. Determine the value of  $n$ .

$$\sum_{c=1}^n (2 - 3c) = -330$$

9. Determine the value of  $n$ :

$$\sum_{k=1}^n (5k - 4) = 2\,205$$

10. Solve the following:

10.1 The common difference of an arithmetic series is 3. Calculate the values of  $n$  for which the  $n^{\text{th}}$  term of the series is 93, and the sum of the first  $n$  terms is 975 .

10.2 Explain why there are two possible answers.

11. The third term of an arithmetic sequence is  $-7$  and the seventh term is 9. Determine the sum of the first 51 terms of the sequence.

12. The second term of an arithmetic sequence is  $-4$  and the sum of the first six terms of the series is 21.

12.1 Find the first term and the common difference.

12.2 Hence determine the value of  $T_{100}$ .

13. The sum of  $n$  terms of an arithmetic series is  $5n^2 - 11n$  for all values of  $n$ . Determine the common difference.

14. The sum of an arithmetic series is 100 times its first term, while the last term is 9 times the first term. Calculate the number of terms in the series if the first term is not equal to zero.

---

15. The first two terms of an arithmetic series are  $m$  and  $n$ .

15.1 Determine the  $5^{\text{th}}$  term in terms of  $m$  and  $n$ .

15.2 Determine the sum of the first 5 terms in terms of  $m$  and  $n$ .

15.3 If  $m = 7$  and  $n = 7$ , determine the sum of the first 5 terms.

16. Solve for  $l$ :  $\sum_{x=1}^4 (3k - l) = 14$

17. Determine the sum of the first 200 numbers where all the multiples of 3 has been removed.

18. The sum to  $n$  terms of an arithmetic series is  $S_n = -\frac{3}{2}n^2 + \frac{7}{2}n$

18.1 Determine the value of the first term of the series.

18.2 Determine the value of the  $6^{\text{th}}$  term of the series.

## 8.6 Exercise 6

1. Prove that

$a + ar + ar^2 + \dots + ar^{n-1} = \frac{a(r^n - 1)}{r - 1}$  and state any restrictions.

2. Solve for  $n$ :  $\sum_{k=1}^n 4(1, 2)^{k-1} < 29\frac{479}{625}$

3. Solve for  $n$ :  $\sum_{k=-2}^n 3\left(\frac{1}{2}\right)^{k+2} > 5\frac{253}{256}$

4. Show that the sum of the first  $n$  terms of the geometric series

$54 + 18 + 6 + \dots + 5\left(\frac{1}{3}\right)^{n-1}$  is given by  $(81 - 3^{4-n})$

5. The eighth term of a geometric sequence is 640. The third term is 20. Find the sum of the first 7 terms.

6. Given:

$\sum_{t=1}^n 8\left(\frac{1}{2}\right)^t$  Calculate the number of terms in the series if  $S_n = 7\frac{63}{64}$

7. The ratio between the sum of the first three terms of a geometric series and the sum of the  $4^{\text{th}}$ ,  $5^{\text{th}}$  and  $6^{\text{th}}$  terms of the same series is  $8 : 27$ .

Determine the constant ratio and the first 2 terms if the third term is 8.

8. If  $x + 1$ ;  $x - 1$ ;  $2x - 5$  are the first three terms of a geometric series. Calculate the value of  $x$

9. Given the geometric series

$(x - 3) + (x - 3)^2 + (x - 3)^3 + \dots$  Write down the value of the common ration in terms of  $x$ .

---

## 8.7 Exercise 7

1. For the general term:  $T_n = 2n$

- Determine if it forms an arithmetic or geometric series.
- Calculate  $S_1, S_2, S_{10}$  and  $S_{100}$
- Determine if the series is convergent or divergent.

2. What value does the series  $\left(\frac{2}{5}\right)^n$  approach as  $n$  tends towards  $\infty$ ?

3. Find the sum to infinity of the following geometric series:

3.1  $3 + 1 + \frac{1}{3} + \frac{1}{9} + \dots$

3.2  $4 + 2 + 1 + \frac{1}{2} + \dots$

3.3  $5 + 1 + \frac{1}{5} + \frac{1}{25} + \dots$

4. Use the sum to infinity to show that:

4.1  $0, \dot{9}$

4.2  $1, \dot{1}\dot{3}$

4.3  $5, \dot{4}$

5. Consider the following:  $\sum_{k=0}^{\infty} \frac{2}{3} \left(\frac{1}{3}\right)^{k-1}$

5.1 Determine the first FOUR terms of the series.

5.2 Determine the sum to infinity of the series.

6. Consider the following:  $\sum_{n=0}^{\infty} 3(0,2)^{n-1}$

6.1 Determine the first FOUR terms of the series.

6.2 Determine the sum to infinity of the series.

7. Determine for which values of  $x$ , the geometric series will converge.

7.1  $2 + \frac{2}{3}(x+1) + \frac{2}{9}(x+1)^2 + \dots$

7.2  $\frac{1}{2} + \frac{1}{2}(x+1) + \frac{1}{2}(x+1)^2 + \dots$

7.3  $3(m-3) + 3(m-2)^2 + 3(m-3)^3 + \dots$

8. The sum to infinity of a geometric series with positive terms is  $4\frac{1}{6}$  and the sum of the first two terms is  $2\frac{2}{3}$ . Find  $a$ , the first term, and  $r$ , the constant ratio between consecutive terms.

9. The sum to infinity of a geometric series is 3 and the constant ratio is  $\frac{1}{2}$ . Find the value of the first term.

10. The first term of an infinite geometric series is 2 and the sum to infinity is 3. Find the value of the common ratio.

---

11.  $2m - 1; m - 2; m - 4$  are the first 3 terms of a geometric series.

11.1 Calculate the values(s) of  $m$ .

11.2 For which values of  $m$  will the series converge?

12. A ball bounces  $\frac{2}{3}$  of its height on each consecutive bounce after it is dropped from a height of 30 m. Calculate the total distance the ball covered.

13. A shrub 110 cm high is planted in a garden. At the end of the first year, the shrub is 120 cm tall. Thereafter the growth of the shrub each year is half of its growth in the previous year. What is the maximum height the shrub will reach?

14. Find  $p$ :

$$\sum_{k=1}^{\infty} 27p^k = \sum_{t=1}^{12} (24 - 3t)$$

## 9 ANSWERS FOR EXERCISES

### 9.1 Exercise 1

1.1  $d = 2 \ 5; 7; 9$

1.2  $d = 11 \ 65; 76; 87$

2.1 constant difference = 4

2.2 26 ; 30 ; 34

2.3  $T_n = 4n - 2$

2.4  $T_{22} = 86$

2.5  $n = 59, 5 \ n$  is not a natural number

3.1 Constant difference =  $-3$

3.2  $-19; -22; -25$

3.3  $T_{20} = -58$

3.4  $T_{32} = -94$

3.5  $n = 24, 33 \ n$  is not a natural number

4.1 Constant difference =  $\frac{1}{2}$

4.2  $\frac{3}{2}; 2; \frac{5}{2}$

4.3  $T_{100} = \frac{95}{2}$  or 47,5

4.4  $T_{68} = \frac{63}{2}$

5.1  $T_5 = 1$

5.2  $T_n = 8,5 - 1,5n$

5.3  $T_{21} = -23$

6.1 Yes, this is an arithmetic sequence since there is a common difference of 4 between consecutive terms.


6.2  $T_{55} = 218$

6.3  $T_{81} = 322$

6.4 This value of  $n$  is not a natural number, therefore 1 204 is not a term of this sequence.

7.1  $T_2 = 3$   $T_3 = 1$   $T_{10} = -13$

7.2



8.1  $x = 7$

---

8.2  $T_{21} = 102$

9.  $x = 3$

10.1  $d = 2b$

10.2  $T_5 = a + 5b$

10.3  $a = 8$   $b = 2$

10.4 4 396

11.  $p = \frac{1}{2}$   $T_{15} = 35\frac{1}{2}$

12. There is a common difference between any two successive terms in the sequence.

The graph of  $T_n$  vs.  $n$  is a straight line.

13. Use the given terms to calculate the common difference

From given terms, we know that  $T_1 = a$

Substitute the values for  $a$  and  $d$  into the equation  $T_n = a + (n-1)d$

Simplify and gather like  $n$  terms.

14.1 4 ; 7 ; 10 ; 13

14.2  $d = 3$

14.3  $T_n = 3n + 1$

14.4  $T_{25} = 76$

14.5  $n = 36$

15.1  $p = 7$

$q = 7$

$r = 35$

$s = 2n + 1$

15.2 801

16.  $T_1 = 6$   $T_2 = 10$   $T_3 = 14$

17.  $T_n = \frac{1}{2}n - 1$

18.  $d = 3$   $T_1 = -10$

19.  $d = -1$

20. 11 terms

21.  $x = 5a - 6$

22.  $a_1 = s$  ;  $a_2 = -s + t$  ;  $a_3 = -3s + 2t$  ;  $a_4 = -5s + 3t$  ;  $a_5 = -7s + 4t$  ;  $a_6 = -9s + 5t$  ;  $a_7 = -11s + 6t$

---

## 9.2 Exercise 2

1.1 Quadratic

1.2 Linear

1.3 Quadratic

1.4 Linear

1.5 Quadratic

1.6 It is neither quadratic nor linear because there is no constant first or second difference.

1.7 Quadratic

1.8 Quadratic

2.  $b = 0$  and  $c = -2$

3.1 Linear

3.2  $T_5 = -7a^2$

3.3  $T_{100} = -197a^2$

4.  $b = 2$  and  $c = -1$

5. 4 ; 14 ; 34 ; 64 ; 104 ; 154

6.1  $d = -1$

6.2  $T_1 = 7$

## 9.3 Exercise 3

1.1  $r = 2$  40 ; 80 ; 160

1.2  $r = \frac{1}{2}$   $\frac{1}{16}$  ;  $\frac{1}{32}$  ;  $\frac{1}{64}$

1.3  $r = 0,1$  0,007 ; 0,0007 ; 0,00007

1.4  $r = 3p$   $27p^4$  ;  $81p^5$  ;  $243p^6$

1.5  $r = -10$  3 000 ; -30 000 ; 300 000

2.1  $T_n = 5(2)^{n-1}$

2.2  $T_n = \left(\frac{1}{2}\right)^n$



---

2.3  $T_n = 7(0,1)^{n-1}$

2.4  $T_n = p(3p)^{n-1}$

2.5  $T_n = -3(-10)^{n-1}$

3.1  $6; 2; \frac{2}{3}$

3.2 Geometric sequence with constant ratio  $r = \frac{1}{3}$

4.  $k = 4$  and  $m = 10$

5.1  $r = 2$

5.2  $a = \frac{1}{4}$

5.3  $T_n = \frac{2^n}{8}$

6.1  $r = 2$  and  $a = \frac{1}{4}$

6.2  $x = \frac{1}{2}$  and  $y = \frac{1}{4}$

6.3  $T_5 = 4$

7.  $T_4 = \sin^3 \theta$  and  $T_5 = \sin^4 \theta$

8.  $x = \frac{1}{4}$  or  $x = 25$   
 $y = \pm \frac{9}{2}$  or  $y = \pm 45$

9.1  $p = 4x^3y$

9.2  $r = \frac{2x}{y}$

10.  $r = \pm \frac{1}{3}$

Therefore possible geometric sequences are:

$-1; \frac{1}{3}; -\frac{1}{9}; \frac{1}{27}; -\frac{1}{81}$  or  $-1; -\frac{1}{3}; -\frac{1}{9}; -\frac{1}{27}; -\frac{1}{81}$

## 9.4 Exercise 4

1.1 8

1.2 5

1.3 34

1.4 2 244

1.5 1

---

2.1 0

2.2 32

2.3  $15a$

2.4 5

3.1 4

3.2  $\frac{15}{16}$

3.3 11

4.1  $\sum_{n=1}^4 (3^{n-3})$

4.2  $\sum_{k=21}^{25} (-1)^k \cdot \frac{1}{k+1}$  or  $\sum_{k=1}^5 (-1)^k \cdot \left(\frac{1}{21+k}\right)$

5.  $\sum_{n=1}^{25} (18 - 7n)$

6.  $\sum_{n=1}^{1000} (2n - 1)$  or  $\sum_{n=0}^{999} (2n + 1)$

7.1  $S_n = \frac{125-5^{3-n}}{4}$

7.2  $S_n = \frac{-3+3\left(-\frac{2}{3}\right)^n}{20}$

## 9.5 Exercise 5

1.1 465

1.2 555

1.3 -60

2.  $S_{300} = 135\,750$

3. 55 195

4.  $\frac{S_{25}}{S_{15}} = \frac{5}{3}$

5.1  $n = 200$

5.2  $S_n = 95\,700$

6.1  $n = 10$

6.2  $T_6 = 46$

7.1  $S_n = 324$

---

7.2  $S_8 + 8 = 316$

8  $n = 15$

9  $n = 30$  or  $n \neq -\frac{147}{5}$

10.1  $n = 13$  or  $n = 50$

10.2 There are two series that satisfy the given parameters:

$$57 + 60 + 63 + \dots + T_{13} = 975$$

$$(-54) + (-51) + (-48) + \dots + T_{50} = 975$$

11.  $S_{51} = 4\,335$

12.1  $a = -9$ ;  $d = 5$

12.2  $T_{100} = 486$

13.  $d = 10$

14.  $n = 20$

15.1  $T_5 = 4n - 3m$

15.2  $S_5 = -5m + 10n$

15.3  $S_5 = -15$

16.  $l = 4$

17. Sum = 13 467

18.1  $S_1 = 2$

18.2  $S_6 = -13$

## 9.6 Exercise 6

1.  $S_n = \frac{a(r^n - 1)}{r - 1}$   
where  $r \neq 1$

2.  $n = 4$

3.  $n = 10$

4.  $(81 - 3^{4-n})$

5. 635

---

6.  $n = 9$

7. Ratio =  $\frac{3}{2}$   
Terms =  $\frac{32}{9}; \frac{16}{3}$

8.  $x = 3 \setminus x = -2$

9. Common ratio =  $(x - 3)$

## 9.7 Exercise 7

1. This is an arithmetic series

$$S_1 = 2$$

$$S_2 = 6$$

$$S_{10} = 110$$

$$S_{100} = 10\,100$$

This is a divergent series

2.  $\frac{2}{3}$

3.1  $S_\infty = \frac{9}{2}$

3.2  $S_\infty = 8$

3.3  $S_\infty = \frac{25}{4}$

4.1  $S_\infty = 1$

4.2  $S_\infty = 1\frac{13}{99}$

4.3  $S_\infty = 5\frac{4}{9}$

5.1  $2 + \frac{2}{3} + \frac{2}{9} + \frac{2}{27}$

5.2  $S_\infty = 3$

6.1  $15 + 3 + \frac{3}{5} + \frac{3}{25}$

6.2  $S_\infty = \frac{75}{4}$

7.1  $-4 < x < 2$

7.2  $-2 < x < 0$

7.3  $2 < m < 4$

8.  $a = \frac{5}{3}$   
 $r = \frac{3}{5}$

---

9.  $a = \frac{3}{2}$

10.  $r = \frac{1}{3}$

11.1  $m = 0$  or  $m = 5$

11.2  $m = 5$

12. 150 m

13. 130 cm

14.  $\frac{2}{3}$